







Pharmacology and Experimental Therapeutics

Investigation of sorbitan niosomal gel for sustained release ocular delivery of piroxicam (1053.3)

Bazigha Abdul Rasool, Oday Alsawad, Hamda Lootah, Iman Abusharbain, and Hiba Abu-Alhaj

Published Online: 1 Apr 2014 **Abstract Number:** 1053.3





Abstract

Objective: In this study, niosomal hydrogel formulation was investigated as a promising carrier for piroxicam sustained release ocular delivery. Methods: Piroxicam loaded niosomes were prepared by thin film hydration technique using various proportions of span and cholesterol. Optimized formulation was selected depending on the highest percent drug entrapment. Morphology, mean size, encapsulation efficiency (EE) and in vitro drug release from niosomes in phosphate buffer pH 7.4, were evaluated. Drugcholesterol interaction was studied by FTIR spectroscopy. Drug retention in niosomes was performed at refrigerated temperature and room temperature for the period of two months. The optimized niosomal formulation was incorporated into 1% carbopol gel and evaluated for the ex vivo transocular drug permeation comparing to a plain drug gel through excised albino rabbit cornea using Franz diffusion cell. Results: Maximum drug EE was obtained with Span 80: cholesterol niosomes of µmolar ratio (300:200). Also they were stable during the storage time. No significant interaction between the drug and cholesterol was detected. The optimized niosomal gel formulation showed prolonged drug release and enhanced drug ocular bioavailability comparing to the plain gel. Conclusion: piroxicam niosomal formulation was prepared successfully and is useful as a sustained release preparation.



Vol. 28, No. 1_supplement

April 2014

Metrics

Downloaded 0 times

Publication History

Published online 1 April 2014



Contact Us

About

Newsletter









9650 Rockville Pike Bethesda, MD 20814 301-634-7000

Terms & Conditions Privacy Policy

© 2019 by the Federation of American Societies for Experimental Biology