

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/289311599>

Theoretically Predicted Descriptors Based Quantitative Structure Activity Relationship Study of the Activity of...

Article in *American Journal of Applied Sciences* · January 2011

CITATIONS

7

READS

5

4 authors:



Bahjat Saeed

University of Basrah

68 PUBLICATIONS 250 CITATIONS

SEE PROFILE



Rita Elias

University of Basrah

22 PUBLICATIONS 52 CITATIONS

SEE PROFILE



Sadiq M-H Ismael

University of Basrah

11 PUBLICATIONS 14 CITATIONS

SEE PROFILE



Kawkab A Hussain

University of Basrah

20 PUBLICATIONS 24 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



isotope effect on C-13 chemical shifts in thio compounds [View project](#)



QSAR studies of flavonoids substituted as anticancer agents. [View project](#)

Theoretically Predicted Descriptors Based Quantitative Structure Activity Relationship Study of the Activity of Acridines Against B-16 Melanoma

¹Bahjat A. Saeed, ²Rita S. Elias, ¹Sadiq M-H. Ismael and ¹Kawkab A. Hussain

¹Department of Chemistry, College of Education,

²Department of Pharmaceutical Chemistry, College of Pharmacy,
University of Basrah, Iraq

Abstract: Problem statement: The probability of success and reducing time and cost in drug discovery process could be increased on the basis of QSAR techniques. The study involves the QSAR investigation of 20 bioactive acridines that have activity against **Approach:** Molecular descriptors, total energy, van der Waals volume, molecular volume, HOMO energy, HOMO-LUMO energy gap, polarizability, refractivity, bond angle of C8-N9-C2 and bond length of C14-N6 were calculated. Initial geometry optimizations were carried out with RM1 Hamiltonian. Lowest energy conformers were subjected to single point calculations by DFT method. Several models for the prediction of biological activity have been drawn up by using the multiple regression technique. **Results:** Four models with R² ranges from 0.88-0.93 were predicted. A model with hepta-parametric equation with R² 0.93 was used to predict the biological activities, the agreement between the observed and the predicted values was up to 93%. **Conclusion:** The biological activity of the studied acridines can be modeled with quantum chemical molecular descriptors.

Key words: Biological activity, single point calculations, quantitative structure, relationship study, quantum chemical, molecular descriptors, biological properties, chemical structures, biological activities, predicted activities

INTRODUCTION

The physiological and biological properties of acridines are well discussed and large number of this kind of compounds have been prepared and evaluated for their biological activities (Su *et al.*, 2006; Cheng *et al.*, 2008). Many of acridine derivatives possess antimicrobial, antiviral and anticancer properties. They are known to be DNA-binding agents as well they interact with other biological targets such as I and II DNA topoisomerase, telomerase, polymerase and protein kinase (Goodell *et al.*, 2009; Kukowska-Kaszuba *et al.*, 2011).

Quantum chemical descriptors have been extensively used in Quantitative Structure-Activity Relationship studies in biochemistry. Numerous reviews have been published on the applications of quantum chemical descriptors (Parthasarathi *et al.*, 2004). The use of quantum chemical descriptors in the development of QSAR has received attention due to reliability and versatility of prediction by these descriptors. For the calculation of the quantum

chemical molecular descriptor used in QSAR studies, semi empirical methods such as AM1 and PM3 mainly have been used (Saeed and Elias, 2010; Saeed *et al.*, 2010a; 2010b). However, DFT method has been used recently for the prediction of physicochemical and biological properties of organic molecules (Shaik *et al.*, 2010; Siu and Che, 2006).

MATERIALS AND METHODS

The studied acridines have been taken with their reactivity from literature (Hansch *et al.*, 2001). Chemical structures and experimental biological activities are gathered in Table 1.

The general formula of the chemical structures of the studied compounds is shown in Fig. 1. Biological activities are presented as log 1/C. All geometries of the acridines are minimized with the semi-empirical RM1 Hamiltonian. Single point calculations have been made at the B3LYP/6-31G level with the RM1 geometry.

Corresponding Author: Bahjat, A. Saeed, Department of Chemistry, College of Education, University of Basrah, Iraq




Fig. 1: The general formula for chemical structures of the studied compounds

Table 1: The chemical structures of the compound studied and their observed activities as log 1/C

Substituents	A _{obs}
4-Cl	2.74
4-CH = CHC ₆ H ₅	2.86
4-CH = C (CN) ₂	3.00
4-CONHC ₃ H ₇	3.25
4-OC ₆ H ₅	3.35
4-NC ₄ H ₉	3.44
4-CN	3.57
3,5-di-NHCONH ₂	3.69
4-Me	3.86
H	4.38
3,4-di-Me	4.45
3,5-di-NH ₂	4.45
3,5-di-OMe	4.52
4-OBu	4.56
4-NHCONH ₂	4.63
4-OMe	4.68
4-(CH ₂) ₃ COOH	4.70
4-OEt	4.87
4-OH	4.93
4-NH ₂	5.35
4-NH ₂ ,3-Me	5.50
4-CH ₂ CH (NH ₂) COOH	6.18

Linear regression analyses are performed to find the best correlation between various biological activity indices and the biological activities of the studied acridines. The values of the descriptors used to develop the Eq. 5 and the predicted activities from this equation are listed in Table 2.

RESULTS AND DISCUSSION

Five models were predicted in this study and have been build up with the use of the following descriptors: van der Waals volume (vWV), polizability (Pol.), energy of HOMO (ϵ_{HOMO}) (eV), total energy (E_T) (hartrees), dipole moment (μ) (deby), C8-N9-C2 bond angle (degrees), C14-N6 bond length (angstroms),

refractivity (Ref.), HOMO-LUMO energy gap ($\Delta\epsilon$) (eV).

The first model is a tetra-parametric equation (Eq. 1) with three descriptors involve: vWV, pol. and ϵ_{HOMO} :

$$1/C = -5.53 \times 10^{-2} (\pm 3.1 \times 10^{-2}) vWV + 0.552 (\pm 0.257) \text{Pol.} + 2.22 (\pm 0.75) \epsilon_{\text{HOMO}} + 21.606 (\pm 2.575) \quad (1)$$

n = 19, R² = 0.83, s = 0.355, F = 24.6 outliers: 4-Cl; 3,5-di-NH₂; 4-CH₂CH(NH₂)COOH

In this model negative values of vWV and Pol. suggest that the activity decreases with an increase in vWV and Pol. of the acridine. While positive value of ϵ_{HOMO} suggests that the activity increases with increasing ϵ_{HOMO} .

The penta-parametric model (Eq. 2) with four descriptors gave better statistical parameters:

$$\text{Log } 1/C = -0.143 \times 10^{-2} (\pm 7.95 \times 10^{-2}) \text{C8-N9-C2} - 5.934 \times 10^{-3} E_T - 0.318 (\pm 7.04 \times 10^{-3}) \mu - 2.49 \times 10^{-2} (\pm 1.30 \times 10^{-3}) vWV + 23.448 (\pm 8.830) \quad (2)$$

n = 19, R² = 0.87, s = 0.294, F = 23.5

outliers: 4-Cl; 4-CH=CH-C₆H₅; 4-CH₂CH(NH₂)COOH

In this model the geometrical parameter C8-N9-C2 bond angle is included and its negative value suggests that better activity could be achieved as the bond angle becomes smaller. This trend is also true for vWV, μ and E_T descriptors. In this case μ has the most important role in predicting the activity.

In Eq. 3 another geometrical parameter, C14-N6 bond length, is included. R² value for this model is of comparable value of that in Eq. 2.

$$\text{Log } 1/C = -5.214 \times 10^{-2} (\pm 7.38 \times 10^{-2}) \text{Ref.} - 0.130 (\pm 0.109) \text{C8-N9-C2} - 25.60 (\pm 43.607) \text{C14-N6} - 0.280 (\pm 9.777 \times 10^{-2}) \mu + 7.898 (\pm 2.113 \times 10^{-2}) vWV - 59.190 (\pm 59.277) \quad (3)$$

n = 20, R² = 0.88, s = 0.317, F = 19.0

outliers: 4-Cl; 4-Me; 4-CH₂CH(NH₂)COOH

In this model the decrease in the bond angle of C8-N9-C2 and bond length of C14-N6 will increase the activity of the acridines. This is also the case with μ and Ref. In contrast an increase in vWV will increase the activity.

The best models concerning the present study the two hepta-parametric equations 4-5. For Eq. 4:

Table 2: The values of the descriptors and the predicted activity (log 1/C) by Eq. 5

	Volume (Å)	μ^a	ref ^b	$\Delta\epsilon^d$	vWV ^c	ϵ_{HOMO}^e	Activity	Observed predicted	Activity ^f Residual
1	1101.2	1.657	122.45	345.70	2.783	-5.000	2.86	2.68	0.18
2	980.7	8.287	105.58	309.57	3.119	-5.568	3.00	3.04	-0.04
3	1045.0	5.521	107.29	329.57	3.358	-5.370	3.25	3.55	-0.30
4	1021.3	3.824	110.78	327.44	3.360	-5.254	3.35	3.37	-0.02
5	1009.9	1.671	105.84	313.30	3.115	-5.280	3.44	3.79	-0.35
6	852.1	7.727	90.28	264.90	3.252	-5.580	3.57	3.40	0.17
7	1054.1	6.300	108.00	333.68	3.290	-4.988	3.69	3.82	-0.13
8	845.7	2.495	89.58	264.30	3.412	-5.189	3.86	4.15	-0.29
9	798.8	2.847	84.54	247.80	3.414	-5.229	4.38	4.12	0.26
10	889.8	2.347	94.60	280.80	3.387	-5.109	4.45	4.39	0.06
11	869.5	3.178	93.90	270.46	3.000	-4.477	4.45	4.35	0.10
12	946.0	4.380	97.47	298.90	3.360	-5.154	4.52	4.72	-0.20
13	1037.5	1.605	104.88	323.80	3.350	-5.077	4.56	4.16	0.40
14	871.0	2.379	91.00	273.35	3.380	-5.060	4.68	4.79	-0.11
15	1027.0	2.418	105.00	325.54	3.396	-5.220	4.70	4.90	-0.20
16	930.6	1.983	95.75	290.10	3.350	-5.088	4.87	4.42	0.45
17	816.9	2.752	86.24	256.00	3.357	-5.061	4.93	4.96	-0.03
18	829.0	0.825	89.24	259.00	3.100	-4.727	5.35	5.59	-0.24
19	875.0	1.046	94.29	275.64	3.090	-4.670	5.5	5.58	-0.08
20	996.0	4.920	103.60	320.84	3.320	-5.270	6.18	5.79	0.39

A: Dipole moment (Deby); b: refractivity; c: van derWals volume; d: $\epsilon_{LUMO} - \epsilon_{HOMO}$ (eV); E: Enegy of HOMO; f: predicted activity expressed by log 1 C⁻¹. A comparison between observed and predicted values of log 1 C⁻¹ for acridines by using Eq. 5 is shown in Fig. 2




Fig. 2: Observed and predicted values of log 1/C for acridines by using Eq. 5

$$\begin{aligned} \text{Log } 1/C &= -0.270 (\pm 0.113) \mu - 62.974 \\ &(\pm 43.695) C_{14-N6} - 0.211 (\pm 9.09 \times 10^{-2}) \\ &\text{Ref.} + 5.318 \times 10^{-2} (\pm 2.65 \times 10^{-2}) \\ &vWV - 2.597 (\pm 1.855) \Delta\epsilon + 0.424 \\ &(\pm 0.688) \epsilon_{HOMO} + 109.1 (\pm 77.66) \end{aligned} \quad (4)$$

n = 20, R² = 0.91, s = 0.284, F = 21.0
outliers: 4-Cl; 4-CH₂CH(NH₂)COOH

According to Eq. C14-N6 is of high significance and as it is the case in Eq. 3 the activity increases with increasing C14-N6. This is also the case with μ , Ref. and $\Delta\epsilon$ although with less degree. Both vWV and ϵ_{HOMO} act in the opposite direction.

Eq. 5 includes the molecular volume instead of C14-N6 as compared to Eq. 4:

$$\begin{aligned} \text{Log } 1/C &= -6.196 \times 10^{-2} (\pm 0.028) V - 0.258 \\ &(\pm 0.138) \mu - 0.400 (\pm 0.154) \text{Ref.} + 0.294 \\ &0.103 vWV - 5.486 (\pm 2.986) \Delta\epsilon - 0.580 \\ &(\pm 1.271) \epsilon_{HOMO} + 30.883 (\pm 9.894) \end{aligned} \quad (5)$$

n = 20, R² = 0.93, s = 0.290, F = 27.2
outliers: 4-Cl; 4-CH₂CH(NH₂)COOH

According to this equation, ϵ_{HOMO} is of major rule and the negative value suggests that biological activity of acridines increases with a decrease in ϵ_{HOMO} . A comparable rule is also clear for $\Delta\epsilon$, while vWV acts in

the opposite direction. The predicted activities of the studied acridines as calculated by Eq. 5 are gathered in Table 2, in addition a comparison between observed and predicted values of log 1/C for acridines used in the development of Eq. 5 is shown in Fig. 2.

The presence of the outliers could be accounted for as a result of the possibility that the molecules may act by different mechanisms or interact with the receptor in different binding modes and due to the intrinsic noise associated with both the original data and methodological aspects involved in the construction of a QSAR model (Hansch *et al.*, 2001).

The values of R^2 for the QSAR models (Eq. 1-5) range from 0.83-0.93 which suggest that these models explain 83-93% of the variance in the data (Polanski *et al.*, 2006). In addition the smaller the value of s and the larger the value of F , the better the QSAR model. The values of s in the Eq. 1-5 range from 0.284-0.355, while the values of F range from 19.0-27.2 which are statistically significant at the 99% level. The values of R^2 , s and F suggest that the QSAR models (Eq. 1-5) are predictive and validate.

CONCLUSION

The study indicated that QSAR of biological activity represented by log 1/C of acridines can be modeled with density functional theory based quantum mechanical molecular descriptors. The regression equations developed in this study can explain 88-93% of the variance in the data. In addition s and F values support the validity of the models.

REFERENCES

- Cheng, M.K., C. Modi, J.C. Cookson, I. Hutchinson and R.A. Heald *et al.*, 2008. Antitumor polycyclic acridines. 20. search for dna quadruplex binding selectivity in a series of 8,13-dimethylquino[4,3,2-kl]acridinium salts: Telomere-targeted agents. *J. Med. Chem.*, 51: 963-975. DOI: 10.1021/jm070587t.
- Goodell, J.R., B. Svensson and D.M. Ferguson, 2009. Spectrophotometric determination and computational evaluation of the rates of hydrolysis of 9-amino-substituted acridines. *J. Chem. Inform. Model.*, 46: 876-883. DOI: 10.1021/ci050322s
- Hansch, C., A. Kurup, R. Garg and H. Gao, 2001. Chem-Bioinformatics and QSAR: A review of QSAR lacking positive hydrophobic terms. *Chem. Rev.*, 101: 619-672. DOI: 10.1021/cr0000067
- Kukowska-Kaszuba, M., K. Dzierzbicka, M. Serocki and A. Skladanowski, 2011. Solid Phase Synthesis and Biological Activity of Tuftsin conjugates. *J. Med. Chem.*, 54: 2447-2454. DOI: 10.1021/jm200002s
- Parthasarathi, R., V. Subramanian, D.R. Roy and P.K. Chattaraj, 2004. Electrophilicity indexes as possible descriptors of biological activity. *Bioorg. Med. Chem.*, 12: 5533-5543. DOI: 10.1061/j.bmc.2004.08.013
- Polanski, J., A. Bak, R. Gieleciak and T. Magdziarz, 2006. Modeling robust QSAR. *J. Chem. Inform. Model.*, 46: 2310-2318. DOI: 10.1021/ci050314b
- Saeed, B.A. and S.R. Elias, 2010. Density functional theory based quantitative structure activity relationship study of 2, 5-bis(1-aziridinyl)-p-benzoquinones with lymphoid leukemia. *Am. J. Appl. Sci.*, 7: 889-892. DOI: 10.3844/ajassp.2010.902.905
- Saeed, B.A. K.Y. Saour, R.S. Elias, N.A. Al-Masoudi and P. La Cola, 2010a. Antitumor and quantitative structure activity relationship study for dihydropyridones derived from curcumin. *Am. J. Immun.*, 6: 7-10. DOI: 10.3844/ajassp.2010.902.905
- Saeed, B.A., K.Y. Saour, R.S. Elias and N.A. Al-Masoudi, 2010b. Antiviral and quantitative structure activity relationship study for dihydropyridones derived from curcumin. *Am. J. Immun.*, 6: 25-28. DOI: 10.3844/ajissp.2010.402.902
- Shaik, S., S. Cohen, Y. Wang, H. Chen, D. Kumar and W. Thiel, 2010. P450-enzymes: Their structures, reactivity and selectivity modeled by QM/MM calculations. *Chem. Rev.*, 110: 949-1017. DOI: 10.1021/cr900121s
- Siu, F.M. and C.M. Che, 2006. Quantitative structure activity (affinity) relationship (QSAR) study on protonation and cationization of α -amino acids. *J. Phys. Chem.*, 110: 12348-12354. DOI: 10.1012/jp064332n
- Su, T.L., Y.W. Lin, T.C. Chou, X. Zhang and V. A. Bacherikov *et al.*, 2006. Potent antitumor 9-anilinoacridines and acridines bearing an alkylating n-mustard residue on the acridine chromophore: synthesis and biological activity. *J. Med. Chem.*, 49: 3710-3718. DOI: 10.1021/jm060197r